

Control of Metazoan Heme Homeostasis by a Conserved Multidrug Resistance Protein

Tamara Korolnek,¹ Jianbing Zhang,¹ Simon Beardsley,¹ George L. Scheffer,² and Iqbal Hamza^{1,*}

¹Department of Animal and Avian Sciences and Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD 20742, USA

²Department of Pathology, VU University Medical Center, 1007 MB Amsterdam, the Netherlands

*Correspondence: hamza@umd.edu

<http://dx.doi.org/10.1016/j.cmet.2014.03.030>

SUMMARY

Several lines of evidence predict that specific pathways must exist in metazoans for the escorted movement of heme, an essential but cytotoxic iron-containing organic ring, within and between cells and tissues, but these pathways remain obscure. In *Caenorhabditis elegans*, embryonic development is inextricably dependent on both maternally derived heme and environmentally acquired heme. Here, we show that the multidrug resistance protein MRP-5/ABCC5 likely acts as a heme exporter, and targeted depletion of *mrp-5* in the intestine causes embryonic lethality. Transient knockdown of *mrp5* in zebrafish leads to morphological defects and failure to hemoglobinize red blood cells. MRP5 resides on the plasma membrane and endosomal compartments and regulates export of cytosolic heme. Together, our genetic studies in worms, yeast, zebrafish, and mammalian cells identify a conserved, physiological role for a multidrug resistance protein in regulating systemic heme homeostasis. We envision other MRP family members may play similar unanticipated physiological roles in animal development.

INTRODUCTION

Heme is almost ubiquitously required by living organisms as a prosthetic group in proteins (Hamza and Dailey, 2012). Heme is synthesized in the mitochondrial matrix but must be trafficked to various subcellular compartments for incorporation into hemoproteins in the cytoplasm, endoplasmic reticulum/Golgi, lysosomes, and peroxisomes (Severance and Hamza, 2009). However, unescorted movement of heme within a cell is inherently hazardous due to the reactivity of free heme. It follows that cells must have specific pathways for the directed movement of heme within and between cells and tissues, but these intra- and intercellular pathways remain poorly defined (Hamza and Dailey, 2012; Severance and Hamza, 2009).

The existence of heme effluxers is all but certain, given that free heme is toxic to cells and must be escorted to various subcellular compartments. One such protein, the major facilitator superfamily member FLVCR1, has been identified (Keel et al.,

2008; Quigley et al., 2004). FLVCR1 null mice lack effective erythropoiesis and die as embryos (Keel et al., 2008). Interestingly, mammalian *Flvcr1* encodes two FLVCR isoforms: FLVCR1a localizes to the plasma membrane, while FLVCR1b localizes to mitochondrial membranes (Chiabrando et al., 2012). The erythropoietic defect observed in FLVCR1 mutant mouse embryos has been attributed to the inability of FLVCR1b to export heme from the mitochondria into the cytosol. However, exactly how cytosolic heme reaches hemoproteins located within subcellular organelles remains undefined (Fleming and Hamza, 2012).

We have exploited *Caenorhabditis elegans* as a genetic model organism because this roundworm is a heme auxotroph (Rao et al., 2005). *C. elegans* is dependent on both maternally derived heme for embryonic development and heme acquired from the diet during larval growth (Rao et al., 2005). Heme is imported into the intestine via the conserved heme permease HRG-1 and its paralog, HRG-4 (Figure 1A) (Rajagopal et al., 2008). The intercellular heme-trafficking protein HRG-3 is secreted from the intestine and carries heme to developing embryos (Chen et al., 2011). HRG-2 is an extraintestinal, heme-binding membrane protein that facilitates heme utilization in the worm hypodermis (Chen et al., 2012). How does intestinal heme, derived from the environment, get delivered to hemoproteins in extraintestinal tissues? Are these intercellular heme transport pathways found in vertebrates? Herein, we show that a multidrug resistance protein MRP-5/ABCC5 likely acts as a cellular heme exporter and is essential for viability in *C. elegans*. This conclusion is supported by our genetic studies in yeast, *C. elegans*, zebrafish, and mammalian cell culture models, which ascribe a physiological role for a multidrug resistance protein in regulating systemic heme homeostasis in metazoa.

RESULTS

HRG-3-Independent Pathway for Heme Transport

Our previous studies implicated HRG-3 in the directed trafficking of heme to extraintestinal tissues, including embryos (Figure 1A). However, *hrg-3* mutant embryos are viable unless subjected to severe maternal heme limitations in utero (Chen et al., 2011). In fact, when worms are grown in the presence of >6 μ M heme, *hrg-3* mRNA is undetectable (Chen et al., 2011). Thus, HRG-3 serves as an inducible mechanism for redirecting heme stores only under heme-limiting conditions. These results would also predict that, in the absence of HRG-3, an alternate pathway exists in *C. elegans*.




Figure 1. *mpr-5* Is an Essential Regulator of *C. elegans* Heme Homeostasis

(A) Current model of heme homeostasis pathways in *C. elegans*. HRG-1/4 proteins import heme into the cytosol of intestinal cells, HRG-3 is secreted from the intestine for heme delivery to other tissues, and HRG-2 is a resident endoplasmic reticulum protein involved in heme utilization within the hypodermis.

(B) Loss of *mpr-5* results in a heme depletion signal that can be rescued by dietary heme. Left: GFP fluorescence (60–120 worms per treatment) quantified using COPAS BioSort in IQ6011 [$P_{hrg-1}::GFP::unc-54\ 3'UTR$; *unc-119(ed3)*; *unc-119* rescue fragment] exposed to vector, *hrg-4*, or *mpr-5* by feeding RNAi at varying heme concentrations. *** $p < 0.001$ when compared to vector control under the same conditions (two-way ANOVA, Bonferroni post test). Right: images of IQ6011 RNAi worms supplemented with 50 μM heme. Error bars represent mean \pm SEM.

(C) Loss of *mpr-5*, and no other *mrp*, results in a heme depletion signal in the heme sensor strain IQ6011. GFP fluorescence (60–120 worms per treatment) quantified from the *hrg-1* transcriptional fusion line (IQ6011) exposed to vector or RNAi against an *mrp* gene at 25 μM heme. GFP was quantified using COPAS BioSort. *** $p < 0.001$ when compared to vector control under the same conditions (one-way ANOVA, Bonferroni post test). Error bars represent mean \pm SEM.

(D) RNAi of FLVCR1 homologs in *C. elegans* does not activate a heme depletion signal in IQ6011. GFP fluorescence in IQ6011 was measured as in Figures 1B and 1C. *** $p < 0.001$ when compared to vector control under the same conditions (one-way ANOVA, Bonferroni post test). Error bars represent mean \pm SEM. See also Figure S1.

We postulated that membrane-bound heme transporters would be suitable candidates for regulating systemic heme homeostasis in the worm and consequently impact the regulation of other heme-responsive genes. By individually depleting 288 heme-responsive genes, which included 41 genes encoding transmembrane-domain containing proteins, we uncovered *mrp-5* (F14F4.5) as a potent regulator of the *C. elegans* transgenic heme sensor strain IQ6011. The IQ6011 strain expresses GFP in the intestine from the heme-responsive *hrg-1* promoter; GFP levels in this strain are inversely correlated with heme levels in the worm (Rajagopal et al., 2008; Sinclair and Hamza, 2010). Depletion of *mrp-5* in IQ6011 by RNAi resulted in significantly greater GFP levels compared to control RNAi, indicating that loss of *mrp-5* results in the animal sensing less heme (Figure 1B; Severance et al., 2010). Importantly, this heme depletion signal could be rescued in a concentration-dependent manner by supplementation with dietary heme (Figure 1B). The increased GFP signal observed by knockdown of the intestinal heme importer

hrg-4 could be completely suppressed with 50 μM heme, while the *mrp-5* RNAi signal persisted even at 500 μM , indicating a far more severe defect (Figure 1B). In addition, microarray analysis and quantitative RT-PCR studies show that *mrp-5* is itself a heme-responsive gene, because its mRNA increased over 3-fold under low heme conditions (Severance et al., 2010). Indeed, in silico analysis of the putative *mrp-5* promoter revealed the presence of a canonical 23 bp heme response element, which we have previously shown is necessary and sufficient to mediate the heme-dependent regulation of *hrg-1* in the worm intestine (Sinclair and Hamza, 2010).

Although the *C. elegans* genome contains eight *mrp* genes (Figure S1, red boxes, available online), *mrp-5* is the only *mrp* family member that significantly alters GFP expression in IQ6011 (Figure 1C) and is the sole *mrp* that is transcriptionally responsive to heme (Severance et al., 2010). Moreover, systematically depleting each of the five FLVCR worm homologs had little or no effect on GFP levels in the heme sensor worm




Figure 2. Worm *mrp-5* Is Essential for Embryonic Development and Larval Growth

(A) MRP-5 membrane topology showing an N-terminal membrane-spanning domain (MSD1) consisting of six TMDs, followed by a cytosolic ATP binding cassette (ABC) domain, a second MSD (MSD2), and a second ABC domain. Segments of MRP-5 that are deleted in the *ok2067* allele are shown in red. (B) Dead progeny of vector control or *mrp-5*(RNAi) worms. ****p* < 0.001 when compared to vector control worms under identical conditions, *n* = 3 (two-way ANOVA, Bonferroni post test). Error bars represent mean ± SEM. (C) Top: the *C. elegans mrp-5* gene contains 20 exons across 7 kb of the X chromosome. *mrp-5* mutants harbor a 1.2 kb deletion (*ok2067*) spanning exons 14 through 17. Bottom: wild-type and *mrp-5* broodmates were grown to gravid adult stage at 200 µM heme. Their F₁ progeny were placed as synchronized L1 larvae on plates seeded with OP50 bacteria with or without 200 µM added heme. Representative images of F₁ worms 4 days posthatching are shown. Scale bar, 20 µM. (D) ZnMP staining in vector control or *mrp-5*(RNAi) worms. Top: worms were exposed to RNAi from L1 to L4 larval stages, pulsed with 60 µM ZnMP for 3 hr, and imaged using confocal microscopy. Bottom: quantification of ZnMP staining (mean ± SEM of ten worms). **p* < 0.05 when compared to control worms (one-way ANOVA, Bonferroni post test). See also Figure S2.

(Figure 1D) (Keel et al., 2008; Lipovich et al., 2002; Quigley et al., 2004).

MRP-5 Is Essential for Embryonic and Larval Development

In worms, *mrp-5* (multidrug resistance protein 5) encodes an ABC transporter of the MRP/ABCC family. *C. elegans* MRP-5 is predicted to include two membrane-spanning domains (MSDs), each containing six transmembrane helices, and two intracellular ATP binding cassette (ABC) domains (Figure 2A) (Borst et al., 2000). Worms were analyzed for growth and developmental phenotypes after RNAi depletion of *mrp-5*. When synchronized larvae (P₀) were fed *mrp-5* RNAi bacteria, they showed no major developmental defects and were able to reach adult gravid stage and lay eggs. However, 80% of the F₁ eggs laid by P₀ worms failed to hatch, and the small number that did hatch arrested as L1 larvae (Figure 2B). This striking embryonic lethal phenotype could be rescued by supplementation of the bacterial food with exogenous heme; 95% of F₁ progeny hatched and

became adults in the presence of 500 µM heme. Together, these results indicate that *mrp-5* is required for embryonic development in *C. elegans*.

The strain VC1599 contains a deletion in *mrp-5(ok2067)* located on the X chromosome, spanning exons 14 through 17 (Figure 2C, top panel; Figure S2A), but is genetically balanced by a marked chromosomal translocation because *mrp-5* mutant worms are embryonic lethal (Edgley et al., 2006). The *mrp-5(ok2067)* deletion removed 176 amino acids, including three transmembrane helices (Figure 2A). Consequently, the predicted topology of the mutant protein will contain only nine transmembrane helices, resulting in a dysfunctional protein with the second ABC domain located on extracellular surface (Figure S2B). Although RT-PCR and sequencing analysis reveals the presence of *mrp-5* mRNA in mutant worms (Figure S2C), depletion of *mrp-5* by RNAi in the *mrp-5(ok2067)* mutants does not enhance or result in additional phenotypes, suggesting that *mrp-5(ok2067)* is likely a null mutation (Figure S2D).

We tested whether supplementary heme could rescue the embryonic lethality of *mrp-5* mutants and found that when VC1599 worms were crossed to wild-type N2 worms to eliminate the balancer, viable F2 homozygous *mrp-5* mutant worms were obtained and easily propagated, but only when worms were grown on food supplemented with >200 μ M heme. When grown on plates containing bacteria with no added heme, *mrp-5(ok2067)* homozygous mutant larva arrested at midlarval stages, indicating that *mrp-5* is required during larval development (Figure 2C, lower panel). To determine if the rescue of *mrp-5* lethality by dietary heme was due to the presence of redundant mechanisms for heme transport, we depleted FLVCR homologs in the *mrp-5(ok2067)* background. Only one, B0416.5, showed a significantly enhanced phenotype in the *mrp-5* mutant worms when depleted (Figure S2E); however, this effect could not be rescued in a dose-dependent manner by exogenous heme and is likely unrelated to heme export from the intestine. It is worth noting that this experiment does not rule out the presence of other low-affinity transporters that can compensate for the loss of *mrp-5* or the possibility that at such high dietary concentrations, heme, which can intercalate into membrane lipids, is traversing the membrane without the assistance of a transporter.

Because MRP-5 is a member of the ABCC/MRP transporter family, members of which function as exporters of lipophilic and organic compounds (Borst et al., 2007; Kos and Ford, 2009), we examined whether MRP-5 was involved in the regulation of heme homeostasis in *C. elegans*. Worms in which *mrp-5* had been depleted showed significantly greater accumulation of zinc mesoporphyrin IX (ZnMP), a fluorescent heme analog, in the intestine compared to control worms (Figure 2D). Notably, VC1599 worms, which are heterozygous for *mrp-5*, exhibited haploinsufficiency phenotypes, because they not only accumulated ZnMP in the intestine but were also resistant to the toxic heme analog gallium protoporphyrin IX, indicating that heme analogs entered the intestine but were poorly accessible to extraintestinal cells (Figures S2F and S2G).

MRP-5 Deficiency Prevents Heme Export from the Intestine

A transcriptional reporter was generated using the putative promoter region of *mrp-5* (3 kb upstream of the ATG start codon) fused to a GFP reporter. *Pmrp-5::GFP* was expressed at all developmental stages, with low levels in the hypodermis and in some neurons, and was consistently highly expressed in the intestine and pharynx (Figure 3A), confirming published intestinal and pharyngeal *in silico* gene expression analysis (Contrino et al., 2012; McGhee et al., 2007).

To delineate the subcellular localization of MRP-5, GFP was fused to the C terminus of MRP-5 and expressed from the intestinal *vha-6* promoter (Oka et al., 2001). In the polarized worm intestinal cells, *Pvha-6::MRP-5::GFP* localized to basolateral membranes and to intracellular compartments, reminiscent of basolateral sorting vesicles in *C. elegans* (Chen et al., 2010) (Figure 3B; Figures S3A and S3B). Similar localization was observed for *MRP-5::GFP* expressed from the endogenous *mrp-5* promoter (data not shown). We next determined if the transgene was capable of rescuing the embryonic lethal phenotype induced by *mrp-5* deficiency. RNAi directed against the 3' UTR of *mrp-5* depleted endogenous *mrp-5*, whereas the *MRP-*

5::GFP transgene, which is expressed with the generic *unc-54* 3' UTR, was left intact, as confirmed visually by GFP fluorescence (data not shown). RNAi against the *mrp-5* 3' UTR resulted in a significant reduction in embryonic lethality in *MRP-5::GFP* transgenic worms, indicating that MRP-5::GFP is a functional protein (Figure 3C).

Because MRP-5 is expressed in multiple tissues, it is conceivable that *mrp-5* depletion in any or all of the tissues may contribute to the embryonic lethal phenotype of *mrp-5* mutant worms. To address the contribution of each tissue to the *mrp-5* phenotype, we utilized tissue-specific RNAi worm strains. Worms carrying the *rde-1* mutation are resistant to RNAi; ectopic expression of *rde-1* from a tissue-specific promoter in the *rde-1* mutant background results in RNAi only in that tissue (Qadota et al., 2007). We depleted *mrp-5* in the VP303 (*rde-1* rescue from the intestinal *nhx-2* promoter), WM118 (*rde-1* rescue from the muscle *myo-3* promoter), and NR222 (*rde-1* rescue from the hypodermal *lin-26* promoter) transgenic worm lines. Depletion of *mrp-5* in the intestine fully recapitulated the F₁ embryonic lethality of whole-animal RNAi, while depletion of *mrp-5* either in the hypodermis or muscle had no effect on F₁ viability (Figure 3D). The lethality caused by RNAi in the VP303 strain could be rescued by supplementation with 500 μ M heme in the diet. Thus, the lethality of *mrp-5* mutants can be attributed to the loss of functional MRP-5 specifically in the intestine, even though *mrp-5* is expressed in extraintestinal tissues. This is further supported by the fact that the *MRP-5::GFP* transgene expressed exclusively in the intestine (*Pvha-6::MRP-5::GFP*) is capable of rescuing the hatching phenotype associated with depletion of endogenous *mrp-5* (Figure 3C).

We next determined whether MRP-5 deficiency phenotypes could be overcome by constitutively expressing HRG-3, the intercellular heme delivery protein, from the intestine. Ectopic expression of *hrg-3* from the intestinal *vha-6* promoter was unable to rescue the embryonic lethality of *mrp-5* RNAi (Figure 3E). This was not due to impaired secretion of HRG-3 from the intestine, because *HRG-3::mCherry* still accumulated in the extraintestinal coelomocytes when *mrp-5* was depleted by RNAi (Figure S3C).

To evaluate the heme status in an extraintestinal tissue when *mrp-5* is depleted, we utilized *Phrg-2::HRG-2::YFP* transgenic worms. The *hrg-2* promoter is active only in the hypodermis, and *Phrg-2::HRG-2::YFP* is induced in the hypodermis when heme levels are limiting in that tissue (Chen et al., 2012). Depletion of *mrp-5* resulted in a striking increase in *HRG-2::YFP* levels, and its expression was not fully suppressed until worms were fed 500 μ M heme (Figure 3F). By contrast, worms in which *hrg-4*, the intestinal heme importer, was depleted did not upregulate *Phrg-2::HRG-2::YFP*, indicating that loss of this particular transporter does not result in limiting heme levels in the hypodermis. Taken together, these results provide strong evidence that MRP-5 is the major intestinal heme exporter.

Mrp5 Is Essential for Erythropoiesis in Zebrafish

Vertebrate ABCC5/MRP5 is ~38% identical to worm MRP-5 with similar overall membrane topology. Within the ABCC/MRP family, the lack of an additional amino-terminal MSD (called MSD0) places ABCC5/MRP5 in a distinct group containing the ABCC4/MRP4, ABCC7 (CFTR), and ABCC12/MRP9 proteins




Figure 3. Worm *mrp-5* Encodes a Putative Intestinal Heme Exporter

(A) GFP expression in IQ5051 [*P_{mrp-5}::GFP::unc-54 3'UTR*; *unc-119(ed3)*; *unc-119* rescue fragment] as determined using confocal microscopy. *mrp-5* is expressed in the hypodermis and some neurons and at higher levels in the pharynx and intestine. P, pharynx; I, intestine; H, hypodermis; E, embryo. Scale bars, 20 μ M.

(B) Transgenic IQ5351 worms [*P_{vha-6}::MRP-5::GFP::unc-54 3'UTR*; *unc-119(ed3)*; *unc-119* rescue fragment] expressing an *mrp-5* translational reporter were imaged by confocal microscopy. Dotted lines indicate apical membrane, dashed lines indicate basolateral membrane, and arrowheads indicate lateral membranes between adjacent intestinal cells. Scale bar, 50 μ M.

(C) The *MRP-5::GFP* fusion gene can rescue the embryonic lethality of *mrp-5* RNAi. RNAi targeting the *mrp-5* ORF causes embryonic lethality in both wild-type N2 and transgenic worms. RNAi against the *mrp-5* 3' UTR results in a less severe embryonic lethal phenotype in N2 worms, but this lethality is significantly rescued by expression of the *MRP-5::GFP* transgene from either the *mrp-5* or the intestinal *vha-6* promoter. *** $p < 0.001$ when compared to wild-type N2 worms under identical conditions, $n = 3$ (two-way ANOVA, Bonferroni post test). Error bars represent mean \pm SEM.

(D) Intestinal RNAi of *mrp-5* recapitulates the embryonic lethality of whole animal *mrp-5* RNAi. Wild-type N2 worms and the tissue-specific RNAi strains were grown on RNAi plates with no added heme. Int, intestinal RNAi; Mus, muscle RNAi; Hyp, hypodermal RNAi; $n = 2$. See Results and Figure S2 for further strain information. Error bars represent mean \pm SEM.

(E) Ectopic expression of *hrg-3* does not rescue the embryonic lethality of whole-animal *mrp-5* RNAi. The experiment was performed as in Figure 1C using wild-type N2 worms and worms ectopically expressing HRG-3 and GFP separated by the SL2 intercistronic sequence [*hrg-3(tm2468)*; *P_{vha-6}::HRG-3::ICS::GFP*, *unc-119(ed3)*; *unc-119* rescue fragment] grown on RNAi plates with no added heme. Error bars represent mean \pm SEM.

(F) Loss of *mrp-5* activates an extraintestinal heme depletion signal. Left: yellow fluorescent protein fluorescence (60–100 worms per treatment) quantified using COPAS BioSort in the *hrg-2* translational fusion line IQ8122 [*P_{hrg-2}::HRG-2::YFP::hrg-2 3'UTR*, *unc-119(ed3)*; *unc-119* rescue fragment] exposed to vector, *hrg-4*, or *mrp-5* RNAi at varying heme concentrations. *** $p < 0.001$, ** $p < 0.01$ when compared to vector control worms (two-way ANOVA, Bonferroni post test). Right: representative images of worms grown at 1 μ M heme from the left panel. Left: error bars represent mean \pm SEM.

See also Figure S3.

(Figure S1, blue box) (Toyoda et al., 2008). Although *C. elegans* contains a single *mrp-5*, the gene has undergone a duplication event, and two *MRP5* paralogs are found in most vertebrates (Figure 4A). Interestingly, the human genome contains three paralogs; in addition to *ABCC5/MRP5* on chromosome 3, *ABCC11/MRP8* and *ABCC12/MRP9* are located in tandem on chromosome 16 (Yabuuchi et al., 2001). Orthologs of *ABCC11/MRP8* can be found in other eutherians, including primates,

dogs, and cows, but are not found in rodent genomes. In all analyzed vertebrate species, the closest homolog of *C. elegans mrp-5* is the vertebrate *ABCC5/MRP5*.

Previous studies with HRG-1 have shown that even though zebrafish and worm HRG-1 proteins are only ~20% identical, they are functional orthologs (Rajagopal et al., 2008). Because zebrafish embryos provide a vertebrate animal model to interrogate hematological changes as a function of aberrant heme




Figure 4. *mrp5* Is Required for Zebrafish Erythropoiesis

(A) Phylogenetic analysis of MRP5/ABCC5 clade in *C. elegans* (orange), zebrafish (red), mice (green), and humans (blue). Sequences were aligned using ClustalW, and a phylogenetic tree was generated using the Neighbor-Joining method in MEGA5. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances are in the units of the number of amino acid substitutions per site.

(B) Lateral view of zebrafish *mrp5* expression by whole-mount in situ hybridization using antisense probe, 4 days postfertilization. Anterior is to the left. Sense probe image is shown to indicate background staining. B, brain; N, neural tube.

(C) Knockdown of zebrafish *mrp5* using morpholinos (MO^{mrp5}) results in severe anemia, as indicated by reduced staining of o-dianisidine-positive red cells, as indicated by black arrowheads.

(D) Knockdown of zebrafish *mrp5* using MO^{mrp5} results in reduced red cell formation. Transgenic embryos expressing GFP from the globin locus control region (LCR-GFP) were injected with control MO or MO^{mrp5}. Left: on day 2 postfertilization, percent GFP-positive RBCs were analyzed by FACS. x and y axes measure GFP and forward scatter, respectively; boxed area indicates gate for RBCs. Right: quantification of morphants shown at left; error bars represent mean \pm SEM. For MO^{mrp5} injection, n = 4. **p < 0.01 for MO^{mrp5} morphants compared to control morphants under identical conditions (one-way ANOVA, Bonferroni post test).

(E) Lateral view of zebrafish *gata1* expression in wild-type, control MO, and MO^{mrp5} morphants by whole-mount in situ hybridization using antisense probe 24 hpf. Anterior is to the left.

(F) Quantification of anemia rescue in zebrafish coinjected with *mrp5* cRNA. ***p < 0.001 for MO^{mrp5} morphants compared to control morphants under identical conditions; n = 4. **p < 0.01 for *mrp5* morphants coinjected with rescue cRNA when compared to *mrp5* morphants with no rescue cRNA under identical conditions; n = 3 (one-way ANOVA, Bonferroni post test). See also Figure S4 and Table S1.

homeostasis (Shafizadeh and Paw, 2004), we analyzed the expression and function of *mrp5/abcc5* in zebrafish. Whole-mount in situ hybridization revealed that *mrp5/abcc5* is widely expressed throughout the embryo, with the greatest expression in the developing central nervous system (Figure 4B; Figure S4).

To knock down *mrp5/abcc5*, we injected fish embryos with morpholinos (MOs) specifically targeted against the ATG start codon of *mrp5/abcc5* (MO^{mrp5}) mRNA. Embryos injected with MO^{mrp5} showed severe anemia with very few o-dianisidine-positive red blood cells (RBCs) compared to embryos injected with control MOs (Figure 4C). MO^{mrp5} morphants also exhibited developmental malformations, including body axis curvature defects and enlarged hearts. The anemia phenotype was reproducible using splice junction MO, which targeted specific exon-

intron junctions (data not shown), resulting in a mutant form of Mrp5 that would be functionally equivalent to mutation at the corresponding position in worm *mrp-5* (Figure 2C; Figure S2A).

To quantify the severe anemia phenotype, we analyzed levels of globin-expressing RBCs in the morphant fish. Transgenic zebrafish expressing GFP from the globin locus control region (LCR-GFP) were injected with control and *mrp5* MOs, and morphant blood was analyzed 2 days postfertilization (dpf) for GFP expression. Fish injected with MO^{mrp5} showed significantly decreased GFP-positive RBCs compared to control MO fish (Figure 4D). Correspondingly, *gata1*, a transcription factor required for primitive erythropoiesis (Paik and Zon, 2010) was robustly expressed in wild-type and control MO embryos, but little or no *gata1* staining was observed in MO^{mrp5} morphants (Figure 4E).




Figure 5. Ectopic MRP5 Expression in Yeast Suggests the Protein Plays a Role in Heme Transport

(A) The *hem1* Δ yeast strain was transformed with indicated vectors, grown for 12 hr without added heme or ALA and then grown for 24 hr under the indicated conditions. Yeast growth was assessed by measuring optical density 600 (OD₆₀₀). Left: uninduced yeast which did not express the transgenes showed no difference in growth after 24 hr in the presence of 250 μM ALA. Right: yeast expressing heme importers HRG-1 and HRG-4 grow significantly better than control yeast. Yeast expressing MRP5 grow significantly worse. ***p < 0.001 compared to vector control after 24 hr; n = 3 (two-way ANOVA, Bonferroni post test). Error bars represent mean \pm SEM.

(B) The *hem1* Δ yeast strain was transformed, grown overnight without added heme or ALA, and spotted on plates supplemented with 0.5 μM heme. Plates were incubated at 30°C for 72 hr.

(C) Heme-dependent β -galactosidase activity. The *hem1* Δ yeast strain was transformed with pCYC1-LacZ as well as empty vector, ScPUG1, CeMRP-5, or hMRP5 and grown with the indicated amount of heme or ALA. Cell lysates were then analyzed for β -galactosidase activity, normalized to vector. ***p < 0.001, **p < 0.01, *p < 0.05 when compared to yeast expressing empty vector under identical conditions; n = 2 (one-way ANOVA, Bonferroni post test). Error bars represent mean \pm SEM.

(D) Heme-dependent ferric reductase activity. The *hem1* Δ *fre1* Δ *fre2* Δ *PGK1-FRE1* yeast strain was transformed with indicated vectors and grown with the indicated amount of heme or ALA. Ferric reductase activity from whole cells was analyzed. **p < 0.01, *p < 0.05 when compared to yeast expressing empty vector under identical conditions; n = 2 (one-way ANOVA, Bonferroni post test). Error bars represent mean \pm SEM.

(E and F) The *hem1* Δ yeast strain expressing (E) CeHRG-4-Flag or CeMRP-5-Flag or (F) CeHRG-4 hemagglutinin (HA) or hMRP5-HA was subjected to indirect immunofluorescence microscopy using anti-Flag or anti-HA antibodies and imaged by confocal microscopy.

The *MO^{mp5}* anemia was indeed due to *Mrp5/Abcc5* deficiency, because coinjecting zebrafish with crRNA encoding *Mrp5/Abcc5* significantly corrected the anemia phenotype (Figure 4F; Table S1). Taken together, these data indicate that *mrp5* is critical for zebrafish erythropoiesis and that MRP5 regulation of systemic heme homeostasis is likely conserved from worms to vertebrates.

MRP5 Is a Putative Heme Transporter

To determine whether MRP5 likely transports heme, we exploited previously established assays in yeast and mammalian cells. *Saccharomyces cerevisiae hem1* mutants are unable to synthesize δ -aminolevulinic acid (ALA), a precursor for heme synthesis, and grow poorly even in the presence of exogenous

heme due to an inefficient heme uptake system (Protchenko et al., 2006, 2008). This inadequate growth of *hem1* Δ can be greatly improved by either expression of a heme importer in the presence of heme or supplementation of ALA to the growth medium (Yuan et al., 2012). When *hem1* Δ yeast express the *C. elegans* heme importer HRG-4 (CeHRG-4), they show significantly improved growth in medium supplemented with 0.5 μM heme when compared to uninduced controls (Figure 5A) (Yuan et al., 2012). However, yeast expressing *C. elegans* MRP-5 or human MRP5 showed significantly reduced growth. Indeed, *hem1* Δ yeast expressing CeMRP-5 or hMRP5 showed a reproducible growth defect in dilution spot assays on agar plates (Figure 5B). The reduced growth of yeast expressing CeMRP-5 or hMRP5 was not due to cell toxicity associated with

overexpression of a large polytopic membrane protein, because growth was restored when cells were cotransformed with the heme importer CeHRG-4 (Figure 5B, bottom three rows).

To assess whether MRP5 expression could alter heme levels in the yeast, we measured the activity of β -galactosidase derived from the *lacZ* reporter under control of the *CYC1* promoter. This promoter is activated by Hap1, a transcription factor positively regulated by cytosolic heme levels (Hon et al., 2003). Yeast expressing CeMRP-5 or hMRP5 showed decreased β -galactosidase activity (Figure 5C), a result consistent with the poor growth phenotype in the spot assay. This result was reproducible in yeast grown in the presence of ALA, indicating that MRP5 can also affect availability of endogenously synthesized heme (Figure 5C, right panels). Interestingly, yeast expressing MRP5 showed lower β -galactosidase activity than cells expressing the yeast heme effluxer Pug1p (Protchenko et al., 2008).

To evaluate heme availability in the yeast secretory compartment, we measured ferric reductase activity, because Fre1p acquires a heme cofactor needed for enzymatic activity in the secretory pathway (Dancis et al., 1990). While Pug1p expression had minimal effect, yeast expressing CeMRP-5 or hMRP5 showed significantly greater ferric reductase activity in the presence of heme and ALA (Figure 5D). Although most MRP transporters expressed in yeast localize to the vacuolar membranes (Paumi et al., 2009), indirect immunofluorescence microscopy localized CeMRP-5 and hMRP5 primarily to intracellular compartments that are distinct from the plasma membrane and the vacuole (Figures 5E and 5F). These results indicate that when expressed in yeast, MRP5 proteins are likely capable of exporting heme from the cytosol into intracellular organelles for delivery to hemoproteins such as Fre1p.

MRP5 Alters Heme Levels in the Secretory Compartment

In mammals, MRP5 is expressed almost ubiquitously (Borst et al., 2007; McAleer et al., 1999; Suzuki et al., 2000), and *Mrp5* knockout mice, previously generated in the FVB genetic background, are viable with no overt phenotypes (de Wolf et al., 2007). To determine how loss of *Mrp5* affected heme homeostasis in a mammalian cell model, we generated mouse embryonic fibroblasts (MEFs) from *Mrp5*^{+/+} and *Mrp5*^{-/-} embryos.

Probing MEF lysates with M₅I-10, a monoclonal antibody generated against the first 38 amino acids of the mouse *Mrp5*, revealed a band of the expected molecular weight by immunoblotting (Figure 6A) (Scheffer et al., 2000). The antibody recognized endogenous *Mrp5* in cell lysates from *Mrp5*^{+/+} MEFs, as well as human MRP5 ectopically expressed in *Mrp5*^{-/-} MEFs. Human MRP5 colocalized with the basolateral membrane marker Na⁺-taurocholate cotransporting polypeptide (NTCP) and not with the apical membrane marker syntaxin 3 (SYN3) in polarized MDCKII cells (Figure 6B). In *Mrp5*^{+/+} MEFs, endogenous *Mrp5* was found in punctuate intracellular vesicles throughout the cytoplasm, with some protein on the cell periphery (Figure 6C). Confocal microscopy studies in *Mrp5*^{-/-} MEFs colocalized MRP5 not only with the plasma membrane (WGA) but also partially with the Golgi (galactosyltransferase) and endosomal recycling organelles (Rab4, Rab5, Rab9, and Rab11) (Figure 6D).

To verify the yeast results, we transfected *Mrp5*^{+/+} and *Mrp5*^{-/-} MEFs with an engineered horseradish peroxidase

(HRP) that was confined to the Golgi with a targeting sequence (White et al., 2013). Because holo-HRP requires heme as a cofactor, HRP activity reflects heme availability in the Golgi compartment (White et al., 2013). When heme-depleted MEFs expressing HRP were supplemented with heme in the growth medium, robust HRP activity was detected in *Mrp5*^{+/+} cells, but not *Mrp5*^{-/-} cells; HRP activity was significantly suppressed by 65% to 80% in *Mrp5*^{-/-} cells (Figure 6E). Together, these results in yeast and mouse cells support a composite model in which MRP5 is a heme exporter that transports heme from the cytosol into the lumen of the secretory pathway (Figure 6F).

DISCUSSION

The nematode *C. elegans* is unable to synthesize heme and therefore is innately dependent on a network of heme sensing, trafficking, and transporting molecules to import environmental heme into the intestine and then export this heme to different tissues and subcellular compartments. In the current study, we show that MRP-5 plays an essential role in *C. elegans* heme homeostasis and that heme is potentially the physiologically relevant substrate of MRP5 across metazoans. For almost two decades, MRP5 has been studied as an exporter of cancer drugs, organic anions, and nucleoside monophosphates, although none of these studies provided direct genetic evidence for a physiological role for MRP5 in growth and development (Borst et al., 2007; Kool et al., 1997; Wijnholds et al., 2000). Our conclusions about MRP-5 function are supported by the following findings: (1) targeted *mrp-5* deficiency in the intestine causes embryonic lethality; (2) MRP-5 primarily localizes to the basolateral plasma membrane, and MRP-5 deficiency results in ZnMP accumulation in the worm intestine; (3) *mrp-5* is expressed during all developmental stages and over a wide range of heme concentrations; and (4) functional heme transport assays in yeast suggest that MRP-5 has the capability to export heme. Altogether, these results suggest that MRP-5 is an important membrane-bound heme exporter in *C. elegans*.

An unanticipated consequence of MRP-5 deficiency in worms is the apparent disconnect between heme levels in the intestine and levels in extraintestinal tissues. In the absence of an intestinal heme exporter, it would be expected that heme will accumulate in the intestine and extraintestinal tissues will be heme-deprived, as seen in the *HRG-2::YFP* reporter strain. However, depletion of *mrp-5* also results in robust expression of the *Phrg-1::GFP* intestinal heme reporter, and this occurs when heme is accumulating within the intestine, a condition when such transporters are not normally expressed. If the *hrg-1* promoter was solely regulated by intestinal heme levels, then we would expect intestinal GFP to be suppressed. It could be that *mrp-5* depletion causes compartmentalization of accumulated heme in the intestine such that this heme can no longer be detected by the *hrg-1* promoter. However, another plausible interpretation of this paradox is that intestinal heme levels are integrated with and regulated by "heme signals" from extraintestinal tissues. That low extraintestinal heme levels activate a depletion signal within a heme-loaded intestine implies the existence of a network for communicating heme status between extraintestinal tissues and their sole source of heme, the intestine. We envisage that cellular heme levels in *C. elegans*, and




Figure 6. MRP5 Localizes to the Secretory Pathway and Alters Heme Levels in This Compartment

(A) Immunoblot analysis of *Mrp5* expression in MEFs generated from *Mrp5*^{+/+} and *Mrp5*^{-/-} FVB mice. Cell lysates were resolved on SDS/PAGE and blotted to nitrocellulose membranes for probing with a monoclonal anti-MRP5 antibody.

(B) MDCKII cells stably expressing human MRP5 were transfected with the basolateral marker NTCP-GFP or the apical marker Syntaxin3-GFP and grown to confluency on transwell filters. Polarization of the monolayer was determined after measuring a transient spike in *trans*-epithelial electrical resistance, which remained above baseline level. Cells were fixed and probed with monoclonal anti-MRP5, followed by Alexa 568-conjugated secondary antibody, and imaged using confocal microscopy. A single confocal section (*xy*) is depicted along with composite stacks in side views (*yz*, *xz*).

(C) Immunohistochemistry of endogenous *Mrp5* in *Mrp5*^{+/+} and *Mrp5*^{-/-} mouse MEFs. *Mrp5* staining was performed as in Figure 6B, using an Alexa 488-conjugated secondary antibody. Scale bar, 20 μM.

(D) Immunolocalization of human MRP5 overexpressed in MEFs by confocal microscopy. WGA is used as a plasma membrane (PM) marker, RFP-GalT is a trans-Golgi (TGN) marker, Rab4YFP marks early endosomes (EE) and the endocytic recycling compartment (ERC), Rab5YFP marks EEs, Rab9YFP marks late endosomes (LE), and the TGN, Rab11YFP marks the ERC and TGN. Scale bar, 20 μM.

(E) Heme-dependent horseradish peroxidase (HRP) activity in *Mrp5*^{+/+} or *Mrp5*^{-/-} MEFs. Cells were transfected with Golgi-HRP and then grown for 24 hr in heme-depleted media plus succinyl acetone (HD+SA) for complete heme depletion. Indicated amounts of heme were added back and cells were incubated for a further 24 hr. Cell lysates were harvested and analyzed for peroxidase activity, which was normalized to peroxidase activity from samples not expressing Golgi-HRP and then to the protein concentration of each sample. ****p* < 0.001 for knockout MEFs when compared to wild-type MEFs under identical conditions; *n* = 3 (two-way ANOVA, Bonferroni post test). Error bars represent mean ± SEM.

(F) Proposed model for heme transport by MRP5. In this composite model, based on results from genetic, biochemical, and localization studies in worm and mammalian systems, MRP5 can localize to the plasma membrane for heme export as well as to the secretory pathway for heme delivery to luminal hemoproteins.

plausibly vertebrates, are regulated not solely by internal heme content (cell autonomous) but also by distally located proteins that signal systemic heme requirements to an intertissue heme trafficking network (cell nonautonomous). This prediction is further supported by our findings that depletion of either a heme exporter (*mrp-5*) or a heme importer (*hrg-1*) produces similar, overlapping phenotypes in worms and zebrafish, i.e., that mobilization of heme in and out of tissues is as important as endogenous heme synthesis (Rajagopal et al., 2008; White et al., 2013; Yuan et al., 2012). Although both heme importers and exporters are obviously essential for survival of a heme

autotroph, these proteins also play an important role in vertebrates as demonstrated by developmental and blood defects in zebrafish.

Given the severe phenotypes associated with *mrp-5* deficiency in worms and zebrafish, why do *Mrp5* null mice not exhibit any overt hematological phenotypes? Clearly, worm and human MRP5 have similar phenotypes in yeast. One plausible explanation could be that in mammals, FLVCR1 isoforms play a prominent role in heme export, while MRP5 performs a more specialized role (Chiabrando et al., 2012; Fleming and Hamza, 2012; Keel et al., 2008). It is notable that worms in which each

of the five FLVCR homologs were depleted exhibited no heme-dependent phenotypes, signifying that, at least in lower metazoans, MRP5 plays a definitive and essential role in heme homeostasis. A second explanation is the influence of the FVB genetic background of the *Mrp5* mutant mouse (de Wolf et al., 2007). Inherently, the FVB strain has high liver and spleen iron content and is therefore ill-suited for studies of systemic iron homeostasis. This is in contrast to the C57BL/6 strain, which has a much lower liver and spleen iron content and, therefore, presents a mouse model that is more sensitive to perturbation in iron metabolism (Wang et al., 2007). For example, mutations in the iron transporter DMT1 are viable in certain mouse strains but become lethal when backcrossed into the C57BL/6 background (M. Fleming, personal communication). Lastly, in vivo compensatory pathways may exist to overcome heme or iron metabolism defects in *Mrp5*^{-/-} mice. It is noteworthy that in humans and other placental mammals, *MRP5/ABCC5* has two recently described paralogs, *ABCC11* and *ABCC12*; mice and zebrafish genomes contain only *ABCC12* (Kruh et al., 2007; Tammur et al., 2001; Yabuuchi et al., 2001). Although it has been reported that *MRP9/ABCC12* also localizes to the secretory pathway in unpolarized mammalian cells (Ono et al., 2007), the functions of *ABCC11* and *ABCC12* are largely unknown.

In worms, yeast, and mammalian cells, MRP5 localizes to both the plasma membrane and intracellular vesicles. The cell-surface localization of MRP5 could be reconciled by the well-studied function of *ABCC* transporters to efflux substrates into the extracellular milieu, consistent with the expected role of exporting heme from the *C. elegans* intestine into the worm's circulatory system. However, MRP5 is also found in the secretory pathway as part of the endocytic recycling compartment. While we do not show that MRP5 has heme export activity by direct biochemical assays or that it can deliver heme to an endogenously expressed mammalian hemoprotein, our genetic and cell biological results in yeast and MEFs support a model in which heme transported into the secretory pathway by MRP5 is incorporated into luminal hemoproteins (Figure 6F). The endosomal trafficking of MRP5 could be mediated by an acidic-dileucine-based sorting signal located in the cytoplasmic carboxy termini of vertebrate MRP5 proteins (Bonifacino and Traub, 2003). It is noteworthy that ATP7A, a copper-transporting P-type ATPase, pumps copper into the secretory compartment for metallation of essential cuproproteins in the Golgi as well as exports copper across the plasma membrane to regulate body copper stores (Lutsenko and Petris, 2003). Conceivably, MRP5 may perform a similar dual function as a heme transporter.

EXPERIMENTAL PROCEDURES

Strains and Vertebrate Experiments

All vertebrate animal experiments were approved by the University of Maryland Institutional Animal Care and Use Committee. Wild-type zebrafish were obtained from the Zebrafish International Resource Center and were staged, raised, and maintained as described previously (Kimmel et al., 1995; Westerfield, 2000). Some worm strains were obtained from the Caenorhabditis Genetics Center, which is funded by National Institutes of Health Office of Research Infrastructure Programs (P40 OD010440). All worm strains used in this study are listed in Table S2. Worms were maintained either in liquid mCeHR2 or on Nematode Growth Medium agar plates (Nass and Hamza, 2007). *C. elegans* transcriptional and translational reporters were generated using Multisite Gateway recombination (Invitrogen) and introduced into

unc-119 worms using the PDS-1000 particle delivery system (Bio-Rad) (Chen et al., 2011).

Worm Sorting and Imaging

Worms were grown from the L1 larval stage to the early adult stage on RNAi plates. Worms for each condition were analyzed for time of flight (length) and extinction (optical density) using a COPAS BioSort (Union Biometrica) with gating parameters for mixed worm populations as in Chen et al. (2011). GFP, yellow fluorescent protein, and ZnMP fluorescence in worms was imaged using a DMIRE2 epifluorescence microscope (Leica) connected to a Retiga 1300 cooled mono 12-bit camera or using a laser scanning confocal microscope (LSM710) (Zeiss).

Zebrafish Experiments

Zebrafish knockdowns were performed using ~1.4 nl per embryo of ~0.5 M MO injected into one-cell-stage embryos (Rajagopal et al., 2008). Embryos were analyzed at 24–72 hr postfertilization (hpf) by *o*-dianisidine staining for hemoglobinization, for LCR-GFP expression, and for *gata1* expression using standard procedures (Ganis et al., 2012). Pools of ~50 embryos were analyzed by fluorescence-activated cell sorting (FACS) from wild-type, control MO, and *mrp5* MO-injected LCR-GFP fish. Whole-mount in situ hybridization was performed on wild-type embryos with an *mrp5* cDNA probe at one cell to 5 dpf, as well as on wild-type, control, and *mrp5* morphants with a *gata1* cDNA probe using standard procedures at 24 hpf (Hauptmann, 1999). Rescue injections were performed using 175 pg of rescue construct.

Yeast Assays

S. cerevisiae strains were grown and assays performed as described previously (Yuan et al., 2012). Plasmids encoding potential heme transporters were transformed into yeast using the lithium method. Before each assay, yeast were heme-starved by growth in 2% w/v raffinose SC (-Ura) liquid medium. Please see Supplemental Experimental Procedures for information regarding the growth assays, the β -galactosidase assay, and the ferric reductase assay.

Mammalian Cell Culture and HRP Assays

Mammalian cell lines were cultured in growth medium consisting of Dulbecco's modified Eagle's medium (DMEM), 10% fetal bovine serum (FBS), and 1% PSG (100 U/ml penicillin, 100 mg/ml streptomycin, and 2 mM L-glutamine). Golgi-targeted HRP was transfected into MEFs using the Lipofectamine transfection reagent (Invitrogen). Following overnight transfection, MEFs were incubated in heme-depleted media (DMEM with 10% heme-depleted FBS and 0.5 mM succinylacetone) for 24 hr. Following heme depletion, cells were switched to heme-depleted media with added heme for 24 hr and harvested for peroxidase activity (White et al., 2013).

Bioinformatics and Statistics

ClustalW and MEGA5 were used to generate a phylogenetic tree for the full-length sequences of all human, mouse, zebrafish, and *C. elegans* MRP/ABCC proteins (Larkin et al., 2007; Tamura et al., 2011). Membrane protein topologies were generated using TMHMM and drawn using TOPO2 (<http://www.sacs.ucsf.edu/TOPO2/>; Krogh et al., 2001). All data are presented as mean \pm SEM. Statistical significance was determined using one-way or two-way ANOVA with Bonferroni post tests in GraphPad Prism, version 5.00 (GraphPad Software, Inc).

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, four figures, and two tables and can be found with this article online at <http://dx.doi.org/10.1016/j.cmet.2014.03.030>.

AUTHOR CONTRIBUTIONS

T.K., S.B., and I.H. designed and executed the *C. elegans* experiments; J.Z. and I.H. designed and executed the zebrafish experiments; T.K. and I.H. designed and executed the mammalian cell experiments; and G.L.S. designed and executed the MRP5 antibody experiment. T.K. and I.H. wrote the manuscript. All authors discussed the results and commented on the manuscript.

ACKNOWLEDGMENTS

We thank Piet Borst, Koen van de Wetering, Harry Dailey, Mike Krause, John Phillips, and Carine White for critical discussions and reading of the manuscript; Piet Borst for the MRP5 cDNA, cell lines, and mice; Barry Paw for the globin LCR-GFP zebrafish strain; Caroline Philpott for the PUG1 plasmid, and John Hanover for use of the COPAS BioSort. This work was supported by funding from the NIH DK85035 and DK74797 (to I.H.) and the Roche Foundation for Anemia Research (to I.H.). I.H. is the CEO and president of Rakta Therapeutics LLC (College Park, MD), a company involved in the development of heme-transporter-related therapeutics.

Received: September 30, 2013

Revised: January 23, 2014

Accepted: March 25, 2014

Published: May 15, 2014

REFERENCES

- Bonifacino, J.S., and Traub, L.M. (2003). Signals for sorting of transmembrane proteins to endosomes and lysosomes. *Annu. Rev. Biochem.* **72**, 395–447.
- Borst, P., Evers, R., Kool, M., and Wijnholds, J. (2000). A family of drug transporters: the multidrug resistance-associated proteins. *J. Natl. Cancer Inst.* **92**, 1295–1302.
- Borst, P., de Wolf, C., and van de Wetering, K. (2007). Multidrug resistance-associated proteins 3, 4, and 5. *Pflugers Arch.* **453**, 661–673.
- Chen, B., Jiang, Y., Zeng, S., Yan, J., Li, X., Zhang, Y., Zou, W., and Wang, X. (2010). Endocytic sorting and recycling require membrane phosphatidylserine asymmetry maintained by TAT-1/CHAT-1. *PLoS Genet.* **6**, e1001235.
- Chen, C., Samuel, T.K., Sinclair, J., Dailey, H.A., and Hamza, I. (2011). An intercellular heme-trafficking protein delivers maternal heme to the embryo during development in *C. elegans*. *Cell* **145**, 720–731.
- Chen, C., Samuel, T.K., Krause, M., Dailey, H.A., and Hamza, I. (2012). Heme utilization in the *Caenorhabditis elegans* hypodermal cells is facilitated by heme-responsive gene-2. *J. Biol. Chem.* **287**, 9601–9612.
- Chiabrando, D., Marro, S., Mercurio, S., Giorgi, C., Petrillo, S., Vinchi, F., Fiorito, V., Fagoonee, S., Camporeale, A., Turco, E., et al. (2012). The mitochondrial heme exporter FLVCR1b mediates erythroid differentiation. *J. Clin. Invest.* **122**, 4569–4579.
- Contrino, S., Smith, R.N., Butano, D., Carr, A., Hu, F., Lyne, R., Rutherford, K., Kalderimis, A., Sullivan, J., Carbon, S., et al. (2012). modMine: flexible access to modENCODE data. *Nucleic Acids Res.* **40** (Database issue), D1082–D1088.
- Dancis, A., Klausner, R.D., Hinnebusch, A.G., and Barriocanal, J.G. (1990). Genetic evidence that ferric reductase is required for iron uptake in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **10**, 2294–2301.
- de Wolf, C.J., Yamaguchi, H., van der Heijden, I., Wielinga, P.R., Hundscheid, S.L., Ono, N., Scheffer, G.L., de Haas, M., Schuetz, J.D., Wijnholds, J., and Borst, P. (2007). cGMP transport by vesicles from human and mouse erythrocytes. *FEBS J.* **274**, 439–450.
- Edgley, M.L., Baillie, D.L., Riddle, D.L., and Rose, A.M. (2006). Genetic balancers. *WormBook Apr* **6**, 1–32.
- Fleming, M.D., and Hamza, I. (2012). Mitochondrial heme: an exit strategy at last. *J. Clin. Invest.* **122**, 4328–4330.
- Ganis, J.J., Hsia, N., Trompouki, E., de Jong, J.L., DiBiase, A., Lambert, J.S., Jia, Z., Sabo, P.J., Weaver, M., Sandstrom, R., et al. (2012). Zebrafish globin switching occurs in two developmental stages and is controlled by the LCR. *Dev. Biol.* **366**, 185–194.
- Hamza, I., and Dailey, H.A. (2012). One ring to rule them all: trafficking of heme and heme synthesis intermediates in the metazoans. *Biochim. Biophys. Acta* **1823**, 1617–1632.
- Hauptmann, G. (1999). Two-color detection of mRNA transcript localizations in fish and fly embryos using alkaline phosphatase and beta-galactosidase conjugated antibodies. *Dev. Genes Evol.* **209**, 317–321.
- Hon, T., Dodd, A., Dirmeier, R., Gorman, N., Sinclair, P.R., Zhang, L., and Poyton, R.O. (2003). A mechanism of oxygen sensing in yeast. Multiple oxygen-responsive steps in the heme biosynthetic pathway affect Hap1 activity. *J. Biol. Chem.* **278**, 50771–50780.
- Keel, S.B., Doty, R.T., Yang, Z., Quigley, J.G., Chen, J., Knoblaugh, S., Kingsley, P.D., De Domenico, I., Vaughn, M.B., Kaplan, J., et al. (2008). A heme export protein is required for red blood cell differentiation and iron homeostasis. *Science* **319**, 825–828.
- Kimmel, C.B., Ballard, W.W., Kimmel, S.R., Ullmann, B., and Schilling, T.F. (1995). Stages of embryonic development of the zebrafish. *Dev. Dyn.* **203**, 253–310.
- Kool, M., de Haas, M., Scheffer, G.L., Scheper, R.J., van Eijk, M.J., Juijn, J.A., Baas, F., and Borst, P. (1997). Analysis of expression of cMOAT (MRP2), MRP3, MRP4, and MRP5, homologues of the multidrug resistance-associated protein gene (MRP1), in human cancer cell lines. *Cancer Res.* **57**, 3537–3547.
- Kos, V., and Ford, R.C. (2009). The ATP-binding cassette family: a structural perspective. *Cell. Mol. Life Sci.* **66**, 3111–3126.
- Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E.L. (2001). Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* **305**, 567–580.
- Kruh, G.D., Guo, Y., Hopper-Borge, E., Belinsky, M.G., and Chen, Z.S. (2007). ABCC10, ABCC11, and ABCC12. *Pflugers Arch.* **453**, 675–684.
- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., et al. (2007). Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947–2948.
- Lipovich, L., Hughes, A.L., King, M.C., Abkowitz, J.L., and Quigley, J.G. (2002). Genomic structure and evolutionary context of the human feline leukemia virus subgroup C receptor (hFLVCR) gene: evidence for block duplications and de novo gene formation within duplicons of the hFLVCR locus. *Gene* **286**, 203–213.
- Lutsenko, S., and Petris, M.J. (2003). Function and regulation of the mammalian copper-transporting ATPases: insights from biochemical and cell biological approaches. *J. Membr. Biol.* **197**, 1–12.
- McAleer, M.A., Breen, M.A., White, N.L., and Matthews, N. (1999). pABC11 (also known as MOAT-C and MRP5), a member of the ABC family of proteins, has anion transporter activity but does not confer multidrug resistance when overexpressed in human embryonic kidney 293 cells. *J. Biol. Chem.* **274**, 23541–23548.
- McGhee, J.D., Sleumer, M.C., Bilenky, M., Wong, K., McKay, S.J., Goszczynski, B., Tian, H., Krich, N.D., Khattri, J., Holt, R.A., et al. (2007). The ELT-2 GATA-factor and the global regulation of transcription in the *C. elegans* intestine. *Dev. Biol.* **302**, 627–645.
- Nass, R., and Hamza, I. (2007). The nematode *C. elegans* as an animal model to explore toxicology in vivo: solid and axenic growth culture conditions and compound exposure parameters. *Curr. Protoc. Toxicol.* **1**, Unit 1.9.
- Oka, T., Toyomura, T., Honjo, K., Wada, Y., and Futai, M. (2001). Four subunit isoforms of *Caenorhabditis elegans* vacuolar H⁺-ATPase. Cell-specific expression during development. *J. Biol. Chem.* **276**, 33079–33085.
- Ono, N., Van der Heijden, I., Scheffer, G.L., Van de Wetering, K., Van Deemter, E., De Haas, M., Boerke, A., Gadella, B.M., De Rooij, D.G., Neeffes, J.J., et al. (2007). Multidrug resistance-associated protein 9 (ABCC12) is present in mouse and boar sperm. *Biochem. J.* **406**, 31–40.
- Paik, E.J., and Zon, L.I. (2010). Hematopoietic development in the zebrafish. *Int. J. Dev. Biol.* **54**, 1127–1137.
- Paumi, C.M., Chuk, M., Snider, J., Stagljar, I., and Michaelis, S. (2009). ABC transporters in *Saccharomyces cerevisiae* and their interactors: new technology advances the biology of the ABC (MRP) subfamily. *Microbiol. Mol. Biol. Rev.* **73**, 577–593.
- Protchenko, O., Rodriguez-Suarez, R., Androphy, R., Bussey, H., and Philpott, C.C. (2006). A screen for genes of heme uptake identifies the FLC family required for import of FAD into the endoplasmic reticulum. *J. Biol. Chem.* **281**, 21445–21457.
- Protchenko, O., Shakoury-Elizeh, M., Keane, P., Storey, J., Androphy, R., and Philpott, C.C. (2008). Role of PUG1 in inducible porphyrin and heme transport in *Saccharomyces cerevisiae*. *Eukaryot. Cell* **7**, 859–871.

- Qadota, H., Inoue, M., Hikita, T., Köppen, M., Hardin, J.D., Amano, M., Moerman, D.G., and Kaibuchi, K. (2007). Establishment of a tissue-specific RNAi system in *C. elegans*. *Gene* 400, 166–173.
- Quigley, J.G., Yang, Z., Worthington, M.T., Phillips, J.D., Sabo, K.M., Sabbath, D.E., Berg, C.L., Sassa, S., Wood, B.L., and Abkowitz, J.L. (2004). Identification of a human heme exporter that is essential for erythropoiesis. *Cell* 118, 757–766.
- Rajagopal, A., Rao, A.U., Amigo, J., Tian, M., Upadhyay, S.K., Hall, C., Uhm, S., Mathew, M.K., Fleming, M.D., Paw, B.H., et al. (2008). Haem homeostasis is regulated by the conserved and concerted functions of HRG-1 proteins. *Nature* 453, 1127–1131.
- Rao, A.U., Carta, L.K., Lesuisse, E., and Hamza, I. (2005). Lack of heme synthesis in a free-living eukaryote. *Proc. Natl. Acad. Sci. USA* 102, 4270–4275.
- Scheffer, G.L., Kool, M., Heijn, M., de Haas, M., Pijnenborg, A.C., Wijnholds, J., van Helvoort, A., de Jong, M.C., Hooijberg, J.H., Mol, C.A., et al. (2000). Specific detection of multidrug resistance proteins MRP1, MRP2, MRP3, MRP5, and MDR3 P-glycoprotein with a panel of monoclonal antibodies. *Cancer Res.* 60, 5269–5277.
- Severance, S., and Hamza, I. (2009). Trafficking of heme and porphyrins in metazoa. *Chem. Rev.* 109, 4596–4616.
- Severance, S., Rajagopal, A., Rao, A.U., Cerqueira, G.C., Mitreva, M., El-Sayed, N.M., Krause, M., and Hamza, I. (2010). Genome-wide analysis reveals novel genes essential for heme homeostasis in *Caenorhabditis elegans*. *PLoS Genet.* 6, e1001044.
- Shafizadeh, E., and Paw, B.H. (2004). Zebrafish as a model of human hematologic disorders. *Curr. Opin. Hematol.* 11, 255–261.
- Sinclair, J., and Hamza, I. (2010). A novel heme-responsive element mediates transcriptional regulation in *Caenorhabditis elegans*. *J. Biol. Chem.* 285, 39536–39543.
- Suzuki, T., Sasaki, H., Kuh, H.J., Agui, M., Tatsumi, Y., Tanabe, S., Terada, M., Saijo, N., and Nishio, K. (2000). Detailed structural analysis on both human MRP5 and mouse *mrp5* transcripts. *Gene* 242, 167–173.
- Tammur, J., Prades, C., Arnould, I., Rzhetsky, A., Hutchinson, A., Adachi, M., Schuetz, J.D., Swoboda, K.J., Ptáček, L.J., Rosier, M., et al. (2001). Two new genes from the human ATP-binding cassette transporter superfamily, ABCC11 and ABCC12, tandemly duplicated on chromosome 16q12. *Gene* 273, 89–96.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., and Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739.
- Toyoda, Y., Hagiya, Y., Adachi, T., Hoshijima, K., Kuo, M.T., and Ishikawa, T. (2008). MRP class of human ATP binding cassette (ABC) transporters: historical background and new research directions. *Xenobiotica* 38, 833–862.
- Wang, F., Paradkar, P.N., Custodio, A.O., McVey Ward, D., Fleming, M.D., Campagna, D., Roberts, K.A., Boyartchuk, V., Dietrich, W.F., Kaplan, J., and Andrews, N.C. (2007). Genetic variation in *Mon1a* affects protein trafficking and modifies macrophage iron loading in mice. *Nat. Genet.* 39, 1025–1032.
- Westerfield, M. (2000). *The Zebrafish Book: A Guide for the Laboratory Use of Zebrafish (Danio rerio)*. (Eugene: Univ. of Oregon Press).
- White, C., Yuan, X., Schmidt, P.J., Bresciani, E., Samuel, T.K., Campagna, D., Hall, C., Bishop, K., Calicchio, M.L., Lapierre, A., et al. (2013). HRG1 is essential for heme transport from the phagolysosome of macrophages during erythrophagocytosis. *Cell Metab.* 17, 261–270.
- Wijnholds, J., Mol, C.A., van Deemter, L., de Haas, M., Scheffer, G.L., Baas, F., Beijnen, J.H., Scheper, R.J., Hatse, S., De Clercq, E., et al. (2000). Multidrug-resistance protein 5 is a multispecific organic anion transporter able to transport nucleotide analogs. *Proc. Natl. Acad. Sci. USA* 97, 7476–7481.
- Yabuuchi, H., Shimizu, H., Takayanagi, S., and Ishikawa, T. (2001). Multiple splicing variants of two new human ATP-binding cassette transporters, ABCC11 and ABCC12. *Biochem. Biophys. Res. Commun.* 288, 933–939.
- Yuan, X., Protchenko, O., Philpott, C.C., and Hamza, I. (2012). Topologically conserved residues direct heme transport in HRG-1-related proteins. *J. Biol. Chem.* 287, 4914–4924.